



MSH483

V1.0 01/2024  
Locally Printed

DO NOT WRITE IN THIS BINDING MARGIN

Queensland  
Government

**Metro South Health**  
**Emergency Department**  
**Opioid Poisoning Nursing Care**  
**Pathway**

**Facility:**

- PAH
- Logan - Beaudesert
- QEH
- Redlands
- Other: .....

(Affix identification label here or complete if E-form)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex:  M  F  I**INTERVENTION****INITIAL**


For patients managed with suspected recreational exposure or deliberate self-poisoning of opioids including heroin, morphine, fentanyl, oxycodone, codeine, buprenorphine, methadone, hydrocodone, tramadol and tapentadol

Ensure patient is on continuous oximetry

Avoid supplemental oxygen unless there is clear evidence of aspiration

Commence vital signs monitoring

- q5min for 15 min following naloxone, then q15min thereafter for 2 hours and q30min for 2 hours (total 4 hours following naloxone)



Time	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
Sats															
RR															
Naloxone	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N
Dose															



Queensland  
Government

**Metro South Health**

**Emergency Department  
Opioid Poisoning Nursing Care  
Pathway**

(Affix identification label here or complete if E-form)

URN:

Family name:

Given name(s):

Address:

Date of birth:

Sex:  M  F  I

Time	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
Sats																
RR																
Naloxone	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N
Dose																
Time	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
Sats																
RR																
Naloxone	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N
Dose																
Time	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
Sats																
RR																
Naloxone	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N	Y / N
Dose																